

Superior Mesenteric Artery Syndrome Treated by Laparoscopic Duodenojejunostomy

El-Khoury E^{1,*}, El-Attrache Najib³ and El-Darazi E²

¹Department of Surgery, Central Military Hospital, Beirut, Lebanon

²Department of Nutrition, Faculty of Arts and Sciences, USEK, Lebanon

³Department of Surgery, Lebanese University, Lebanon

Volume 4 Issue 1- 2020

Received Date: 29 Apr 2020

Accepted Date: 08 May 2020

Published Date: 11 May 2020

1. Abstract

Superior Mesenteric Artery syndrome (SMA sd) is a very rare disease. It is suspected in the case of intestinal obstruction in severely underweight patients. When conservative management fails surgery becomes needed, with laparoscopic duodenojejunostomy being the best approach.

2. Keywords:

Superior Mesenteric Artery; Duo-denajejunostomy; Laparoscopy

3. Case report

A 22-year-old patient, with a BMI of 22 kg/m², presented to us with a history of intestinal obstruction for more than five days, and a severely distended abdomen on physical exam. Abdominal CT Scan confirmed the diagnosis of SMA sd with an aortomesenteric angle of 20 degrees and a distance of 7 mm (Figure 1). He was managed conservatively with nasojejunal feeding, with no relief in his symptoms after five days. Laparoscopic duodenojejunostomy was decided. Under general anesthesia, the patient was positioned in Trendelenbourg decubitus positon and the surgeon stood between his legs. The Camera trocar was placed 2 cm below the umbilicus and the two operating trocars at the midclavicular line bilaterally on the same level. The transverse colon and mesocolon were retracted superiorly till clear vision of the third part of the duodenum (D3) (Figure 2-3). The retroperitoneum is entered by incising the thin peritoneal layer below the middle of D3, the Inferior Vena Cava (IVC) became visualized. (Figure 4) Then, D3 was completely mobilised from the aorta and the IVC posteriorly (Figure 5), and dissected from its junction with the second duodenum (D2) medially (Figure 6), from the Superior Mesenteric Vein (SMV) laterally and from the transverse mesocolon anteriorly (Figure 7). Finally, the proximal Jejunum was anastomosed mechanically to D3 laterolaterally (Figure 8). The post operative course was uneventful. The patient was relieved from his obstructive symptoms and subsequently regained weight.




Figure 1: Aortomesenteric angle of 20 degrees compressing the distal D3 (arrow). Severely distended stomach (Long arrow)



Figure 2: Third duodenum (Arrow) Retracted transverse colon with its mesentery (grasper)

*Corresponding Author (s): El-Khoury Elias, Department of Surgery, Central Military Hospital, Beirut, Lebanon, E-mail : efkhoury@gmail.com

<http://acmcasereports.com/>

Citation: El-Khoury Elias, El-Darazi Elham and El-Attrache Najib. Superior Mesenteric Artery Syndrome Treated by Laparoscopic Duodenojejunostomy. Annals of Clinical and Medical Case Reports. 2020; 4(1): 1-4.




Figure 3: Retraction of D3 (by the grasper) with identification of its inferior limit (arrow)




Figure 7: Upper part of D3 (arrow) dissected from the transverse mesocolon (longer arrow)




Figure 4: Incision of the peritoneal layer inferior to D3 and entering the retroperitoneum with identification of the IVC (arrow)



Figure 8: Mechanical duodenojejunostomy

4. Discussion

SMA Sd was first reported by Von Rokitansky on an autopsy [1]. Wilkie further detailed its pathophysiology and diagnostic findings [2]. It is very rare with an incidence below 0.3%, [3, 4] affecting mainly young adults and more commonly female patients [5]. SMA Sd should be on the top of the differential diagnosis in severely underweight patients with signs and symptoms of intestinal obstruction. In normal individuals, the angle created by the SMA and the aorta is around 38 and 56 degrees while the normal distance is 10-20 mm [6, 7]. In SMA Sd the distal part of D3 becomes compressed between the SMA and the aorta when the aortomesenteric angle is less than 25% and the distance is shortened to < 8 mm [8-11]. This is due in the majority of cases to the loss of the mesenteric fat pad, [12] like in extreme weight loss, in some cases in bariatric surgery, cancer, burns, and anorexia nervosa [13]. In a minority of cases, SMA Sd is caused by anatomic variants like short ligament of Treitz, high insertion of duodenum at the ligament of Treitz, low origin of SMA, lumbar hyperlordosis, where the aortomesenteric angle is anatomically more acute or D3 is more cephalad than usual and thus compression is more likely [14]. SMA Sd can be a surgical




Figure 5: Posterior mobilisation of D3 from the IVC (longer arrow) and aorta (arrow)




Figure 6: Medial mobilisation of D3 (arrow) and identification of D2 (longer arrow)

complication like in scoliosis surgery, [14, 15] ileoanal pouch surgery, [16] abdominal aortic aneurysm repair [15]. SMA Sd may rarely coexist with the Nutcracker syndrome when the left renal vein becomes compressed [17, 18]. CT scan is the standard diagnostic radiological tool as it allows the measurement of the aortomesenteric angle and distance. SMA Sd presents like any intestinal obstructive disease. Relief occurs when the patient goes from supine to prone position because gravity contributes to the duodenal compression. Initially, SMA Sd should be treated conservatively, and surgery is needed when medical treatment fails [19, 20]. The surgical options include duodenojejunostomy, gastrojejunostomy and duodenal derotation. Gastrojejunostomy does not relieve the proximal duodenal obstruction and patients can continue to have postoperative vomiting [21]. Duodenal derotation has the advantages that it does not require an anastomosis, however it has a high failure rate [9, 22, 23]. Laparoscopic duodenojejunostomy, first performed in 1998, [24] is considered to be the best surgical treatment for SMA Sd with the same outcome as open surgery but with less morbidity with 90% success rate [6, 17, 22, 25-28].

References

- Von Rokitansky C. In: Lehrbuch der pathologischen Anatomie. 3rd ed. Vienna: Braumüller; 1861. Superior mesenteric artery syndrome; p. 87.
- Wilkie DPD. Chronic Duodenal Ileus. The American Journal of the Medical Sciences. 1927; 173: 643-48.
- Ganss A, Rampado S, Savarino E, Bardini R. Superior Mesenteric Artery Syndrome: a Prospective Study in a Single Institution. *J Gastrointest Surg.* 2019; 23: 997-1005.
- Shiu JR, Chao HC, Luo CC, Lai MW, Kong MS, Chen SY, et al. Clinical and nutritional outcomes in children with idiopathic superior mesenteric artery syndrome. *J Pediatr Gastroenterol Nutr.* 2010; 51: 177-82.
- Smith BM, Zyromski NJ, Purtill MA. Superior mesenteric artery syndrome: an underrecognized entity in the trauma population. *J Trauma.* 2008; 64: 827-30.
- Valiathan G, Wani M, Lanker J, Reddy PK. A Case Series on Superior Mesenteric Artery Syndrome Surgical Management, Single Institution Experience. *J Clin Diagn Res.* 2017; 11: 1-3.
- Ozkurt H, Cenker MM, Bas N, Erturk SM, Basak M. Measurement of the distance and angle between the aorta and superior mesenteric artery: normal values in different BMI categories. *Surg Radiol Anat.* 2007; 29: 595-9.
- Neri S, Signorelli SS, Mondati E, Pulvirenti D, Campanile E, Di Pino L, et al. Ultrasound imaging in diagnosis of superior mesenteric artery syndrome. *J Intern Med.* 2005; 257: 346-51.
- Merrett ND, Wilson RB, Cosman P, Biankin AV. Superior mesenteric artery syndrome: diagnosis and treatment strategies. *J Gastrointest Surg.* 2009; 13: 287-92.
- Oliva-Fonte C, Fernández Rey C, Pereda Rodríguez J, González-Fernández AM. Wilkie's syndrome. *Rev Esp Enferm Dig.* 2017; 109: 62-3.
- Cardarelli-Leite L, Velloni FG, Salvadori PS, Lemos MD, D'Ippolito G. Abdominal vascular syndromes: characteristic imaging findings. *Radiol Bras.* 2016; 49: 257-63.
- Biank V, Werlin S. Superior mesenteric artery syndrome in children: a 20-year experience. *J Pediatr Gastroenterol Nutr.* 2006; 42: 522-5.
- Mascolo M, Dee E, Townsend R, Brinton JT, Mehler PS. Severe gastric dilatation due to superior mesenteric artery syndrome in anorexia nervosa. *Int J Eat Disord.* 2015; 48: 532-4.
- Iwaoka Y, Yamada M, Takehira Y, Hanajima K, Nakamura T, Murohisa G, et al. Superior mesenteric artery syndrome in identical twin brothers. *Intern Med.* 2001; 40: 713-5.
- Zaraket V, Deeb L. Wilkie's Syndrome or Superior Mesenteric Artery Syndrome: Fact or Fantasy? *Case Rep Gastroenterol.* 2015; 9: 194-9.
- Matheus Cde O, Waisberg J, Zewer MH, Godoy AC. Syndrome of duodenal compression by the superior mesenteric artery following restorative proctocolectomy: a case report and review of literature. *Sao Paulo Med J.* 2005; 123: 151-3.
- Shi Y, Shi G, Li Z, Chen Y, Tang S, Huang W. Superior mesenteric artery syndrome coexists with Nutcracker syndrome in a female: a case report. *BMC Gastroenterol.* 2019; 19: 15.
- Barsoum MK, Shepherd RF, Welch TJ. Patient with both Wilkie syndrome and nutcracker syndrome. *Vasc Med.* 2008; 13: 247-50.
- Rodriguez EJ, Hernandez-Villegas AC, Serralde-Zuniga AE, Reyes-Ramirez A. The two sides of superior mesenteric artery syndrome treatment: conservative or surgical management? *Nutr Hosp.* 2017; 34: 997-1000.
- Bohanon FJ, Nunez Lopez O, Graham BM, Griffin LW, Radhakrishnan RS. A Case Series of Laparoscopic Duodenojejunostomy for the Treatment of Pediatric Superior Mesenteric Artery Syndrome. *Int J Surg Res.* 2016; 2016:1-5.
- Pottorf BJ, Husain FA, Hollis HW Jr, Lin E. Laparoscopic management of duodenal obstruction resulting from superior mesenteric artery syndrome. *JAMA Surg.* 2014; 149: 1319-22.

22. Lorentziadis ML. Wilke's syndrome: A rare cause of duodenal obstruction. *Ann Gastroenterol.* 2011; 24: 59-61.
23. Reece K, Day R, Welch J. Superior Mesenteric Artery Syndrome with Abdominal Compartment Syndrome. *Case Rep Emerg Med.* 2016; 2016: 7809281.
24. Gersin KS, Heniford BT. Laparoscopic duodenojejunostomy for treatment of superior mesenteric artery syndrome. *JSLS.* 1998; 2: 281-4.
25. Wang T, Wang ZX, Wang HJ. Clinical Insights into Superior Mesenteric Artery Syndrome with Multiple Diseases: A Case Report. *Dig. Dis. Sci.* 2019; 64: 1711-4.
26. Kirby GC, Faulconer ER, Robinson SJ, Perry A, Downing R. Superior mesenteric artery syndrome: a single Centre experience of laparoscopic duodenojejunostomy as the operation of choice. *Ann R Coll Surg Engl.* 2017; 99: 472-5.
27. Wyten R, Kelty CJ, Falk GL. Laparoscopic duodenojejunostomy for the treatment of superior mesenteric artery (SMA) Syndrome: case series. *J Laparoendosc Adv Surg Tech A.* 2010; 20: 173-6.
28. Barchi LC, Alves AM, Jacob CE, Bresciani CJC, Yagi OK, Nogueira TG, et al. Favorable minimal invasive surgery in the treatment of superior mesenteric artery syndrome: Case report. *Int J Surg Case Rep.* 2016; 29: 223-6.