

Renal Failure and Quality of Life Indicators in Kidney Transplantation

Maria Dalamagka*

Anesthesia Department, General Hospital of Larisa, Greece

*Corresponding author:

Maria Dalamagka,
Anesthesia Department, General Hospital of Larisa,
Greece, E-mail: mary.dalamaga@gmail.com

Received: 10 Apr 2022

Accepted: 25 Apr 2022

Published: 30 Apr 2022

J Short Name: ACMCR

Copyright:

©2022 Maria Dalamagka. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially.

Citation:

Maria Dalamagka, Renal failure and Quality of Life Indicators in Kidney Transplantation. *Ann Clin Med Case Rep.* 2022; V9(2): 1-2

1. Abstract

Health-related quality of life (HRQL) contains many aspects of patients' health such as physical, psychological, social functioning and general well-being. Progress in renal transplantation and immunosuppressive therapies have increased significantly in recent decades, resulting in allograft survival rates at one year is now over 90%. Numerous clinical trials have established the importance of quality of life in a variety of diseases, and it is extremely popular to evaluate quality of life in clinical trials as a measure of patients' subjective state of health. The purpose of the study was to identify factors associated with quality of life after renal transplantation.

2. Introduction

Kidney transplantation is the treatment of choice in end-stage renal failure. The transplantation may be the focus for increasing survival and to maximize quality of life. However, there are certain factors that may affect the quality of life after transplantation, such as side effects from highly immunosuppressive drugs, the presence of common disease states and the possibility of rejection. The main goal of transplantation is to achieve maximum quality and longevity while minimizing the impact of disease and health care costs. HRQL is also progressively being recognized as an important outcome measure after organ transplantation. Along with other indicators related to patient improvement and graft survival, quality of life has been assessed as a valid outcome measure. Research on quality of life aims to lead to a broader view of subjective health, as they consider health to be a puzzle of general well-being. It is generally accepted that patients with a functional kidney transplant have an improved quality of life compared to patients undergoing dialysis. Specific tools in assessing the quality of life in a

kidney transplant are: the kidney transplant questionnaire (KTQ), the quality of life in kidney disease (KDQOL), and the transplant unit's end-stage renal disease (ESRDSC-TM) checklist. The ESRDSC-TM was developed specifically to evaluate the effects of immunosuppressive drugs on quality of life. The authors screened more than 400 transplant patients and evaluated the re-screening correlation in a subset of 88 patients over a one-year period and found sufficient validity. General tools are used for comparisons between groups and studies and to assess the impact of different diseases on quality of life (QOL). These tools are used in research and are as follows: Sickness Impact Profile (SIP), the 36-item modified medical Outcomes Survey (SF-36), and the Nottingham Health Profile (NHP). With more than 200 publications, the SF-36 is one of the most widely used tools for assessing quality of life [1-16].

3. Results

There is evidence for the use of the SF-36 in patients with chronic renal failure. The EQ-5D is favored among the preferred measurements, as there is more evidence, but this ratio is applicable to the above. Since multidimensional specific renal disease measurements in KDQOL include most evidence. Given this overlap between the SF-36 and KDQOL, there is some benefit when used in the same survey, unlike the combination of EQ-5D and KDQOL provides additional information about the perception of patients for kidney disease. While the idea of using a short questionnaire based on severity would be very good, the main benefit would only control or recognition of symptoms. The SF-36 is the only general measurement with good properties and functional characteristics. Further more psychometric criteria can be reproduced when administered as autonomous as possible, and when used in conjunc-

tion with KDQOL questionnaire. The EQ-5D appears to have a favorable use, since three of the four studies that used the EQ-5D were conducted in the UK. The evidence shows high responder rates.

4. Conclusion

In general, quality of life improved after a successful kidney transplant compared with dialysis patients, and the effect was more pronounced in men than in women. These studies clearly show that kidney transplantation is not only the cheapest long-term replacement therapy, but is also associated with lower mortality and better quality of life for patients.

References

1. Bass EB, Wills S, Fink NE. How strong are patients' preferences in choice between dialysis modalities and doses? *Am J Kidney Dis.* 2004; 44: 695-705.
2. Wasserfallen JB, Halabi G, Saudan P. Quality of life on chronic dialysis: comparison between haemodialysis and peritoneal dialysis. *Nephrol Dial Transplant.* 2004; 19: 1594-9
3. Forsberg A, Lorenzon U, Nilsson F, Backmana L. Pain and health related quality of life after heart, kidney, and liver transplantation. *Clin Transplant* 1999; 13: 453-60.
4. Hays RD, Kallich JD, Mapes DL. Development of the kidney disease quality of life (KDQOL) instrument. *Qual Life Res.* 1994; 3: 329-38.
5. Canadian Erythropoietin Study Group. Association between recombinant human erythropoietin and quality of life and exercise capacity of patients receiving haemodialysis. *BMJ* 1990; 300: 573-8
6. Churchill DN, Wallace JE, Ludwin D. A comparison of evaluative indices of quality of life and cognitive function in hemodialysis patients. *Control Clin Trials* 1991; 12(Suppl.): S159-67.
7. Harris DC, Chapman JR, Stewart JH. Low dose erythropoietin in maintenance haemodialysis: improvement in quality of life and reduction in true cost of haemodialysis. *Aust N Z J Med.* 1991; 21: 693-700.
8. Laupacis A, Keown P, Pus N. A study of the quality of life and cost-utility of renal transplantation. *Kidney Int* 1996; 50: 235-42.
9. Sesso R, Yoshihiro MM, Ajzen H. Late diagnosis of chronic renal failure and the quality of life during dialysis treatment. *Braz J Med Biol Res.* 1996; 29: 1283-9.
10. Sesso R, Yoshihiro MM. Time of diagnosis of chronic renal failure and assessment of quality of life in haemodialysis patients. *Nephrol Dial Transplant.* 1997; 12: 2111-6.
11. Cleemput I, Kesteloot K, De Geest S. Health professionals' perceptions of health status after renal transplantation: a comparison with transplantation candidates' expectations. *Transplantation.* 2003; 76: 176-82.
12. Greiner W, Obermann K, Graf vd. Schulenburg J-M. Socioeconomic evaluation of kidney transplantation in Germany. *Arch Hell Med.* 2001; 18: 147-55.
13. Moons P, Vanrenterghem Y, Van Hooff JP. Health related quality of life and symptom experience in tacrolimus-based regimen after renal transplantation: a multicentre study. *Transpl Int.* 2003; 16: 653-64.
14. Polsky D, Weinfurt KP, Kaplan B. A economic and quality-of-life assessment of basiliximab vs antithymocyte globulin immunoprophylaxis in renal transplantation. *Nephrol Dial Transplant.* 2001; 16: 1028-33.
15. Roderick P, Nicholson T, Armitage A. An evaluation of the costs, effectiveness and quality of renal replacement therapy provision in renal satellite units in England and Wales. *Health Technol Assess.* 2005; 9: 1-178.
16. McFarlane PA, Bayoumi AM, Pierratos A, Redelmeier DA. The quality of life and cost utility of home nocturnal and conventional in-center hemodialysis. *Kidney Int.* 2003; 64: 1004-11.